


#11
09/863795

Rudolf Meyer

explosives

2nd, revised and extended edition



Weinheim · Deerfield Beach, Florida · Basel · 1981

Dr. Rudolf Meyer
Trappenbergstraße 2
D-4300 Essen-Stadtwald

Copy Editor: Dr. Gerd Giesler

Deutsche Bibliothek Cataloguing-in-Publication Data

Meyer, Rudolf:

Explosives / Rudolf Meyer. [Sponsored by
WASAG-Chemie]. — 2., rev. and extended ed. —
Weinheim; Deerfield Beach, Florida; Basel:
Verlag Chemie, 1981.

Einheitssacht.: Explosivstoffe -engl.-
ISBN 3-527-25933-3

© Verlag Chemie GmbH, D-6940 Weinheim, 1981

All rights reserved (including those of translation into foreign languages). No part of this book may be reproduced in any form — by photoprint, microfilm, or any other means — nor transmitted or translated into a machine language without written permission from the publishers. Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.


Printed by: Gebr. Diesbach, D-6940 Weinheim
Bookbinder: Georg Krändt, D-6148 Heppenheim
Printed in the Federal Republic of Germany

It is prepared by condensation of formaldehyde with nitromethane and by nitration of the nitroisobutylglycerin product under the same conditions as nitroglycerin. The nitration and stabilization procedures are very difficult because of decomposition reactions.

While being of interest to the explosives industry, since it has an ideal oxygen balance, its stabilization in actual practice proved to be impossible.

Nitromethane

Nitromethan; nitrométhane; NM



colorless liquid

molecular weight: 61.0

energy of formation: $-413.7 \text{ kcal/kg} = -1731 \text{ kJ/kg}$

enthalpy of formation: $-443.1 \text{ kcal/kg} = -1854 \text{ kJ/kg}$

oxygen balance: -39.3%

nitrogen percentage: 22.96 %

volume of detonation gases: 1102 l/kg

heat of explosion (H_2O liq.): $138 \text{ kcal/kg} = 4763 \text{ kJ/kg}$

specific energy: $130 \text{ mt/kg} = 1275 \text{ kJ/kg}$

density: 1.1385 g/cm^3

solidification point: $-29^\circ\text{C} = -20^\circ\text{F}$

boiling point: $101.2^\circ\text{C} = 214.2^\circ\text{F}$

evaporation enthalpy: $151 \text{ kcal/kg} = 631 \text{ kJ/kg}$

vapor pressure:

millibar	temperature $^\circ\text{C}$	$^\circ\text{F}$
1.3	- 29	- 20 (solidification point)
10	0	32
32	20	68
140	50	122
283	80	176
1010	101.2	214.2 (boiling point)

lead block test: $400 \text{ cm}^3/10 \text{ g}$

detonation velocity, confined:

$6290 \text{ m/s} = 20\,700 \text{ ft/s}$ at $\rho = 1.138 \text{ g/cm}^3$

Nitromethane is sparingly soluble in water. The compound is of interest as a solvent rather than as explosive. Its technical synthesis


involves nitration of methane with nitric acid above 400 °C (750 °F) in the vapor phase.

It was used in the USA for underground model explosions ("Pre-Gondola"), in preparation for the → *Nuclear Charge* technique. Nitromethane was also employed in stimulation blasting in oil and gas wells. PLX (*Picatinny Liquid Explosive*) is a mixture of nitromethane with 5 % ethylene diamine and is foreseen to clean up mine fields.

Nitromethane is of interest both as a monergolic and as a liquid fuel for rockets.

Nitromethylpropanediol Dinitrate

methylnitropropanediol dinitrate; Nitromethylpropandiol-dinitrat; dinitrate de nitromethylpropanediol



gross formula: $\text{C}_4\text{H}_7\text{N}_3\text{O}_8$

molecular weight: 225.1

oxygen balance: -24.9 %

nitrogen percentage: 18.67 %

volume of detonation gases: 907 l/kg

heat of explosion (H_2O liq.): 1283 kcal/kg = 5372 kJ/kg

specific energy: 129 mt/kg = 1260 kJ/kg

The product is prepared by condensation of → nitroethane with formaldehyde, after which the nitromethylpropane diol is nitrated.

Nitroparaffins

are aliphatic hydrocarbons with NO_2 -groups attached directly to carbon atoms. They are mainly obtained by nitration in a gaseous state; → *Nitromethane; Nitroethane, Trinitromethane; Tetra-nitromethane*.

Nitroparaffins can be reacted with formaldehyde to obtain nitroalcohols, which can be further esterified with nitric acid (→ e.g., *Nitroisobutylglycerol Trinitrate*).